

# Problem Set AB due May 1, 2017

## MC - Non Calc

1. If  $y = (x^3 + 1)^2$ , then  $\frac{dy}{dx} =$

- (A)  $(3x^2)^2$       (B)  $2(x^3 + 1)$       (C)  $2(3x^2 + 1)$       (D)  $3x^2(x^3 + 1)$       (E)  $6x^2(x^3 + 1)$

2.  $\int_0^1 e^{-4x} dx =$

- (A)  $\frac{-e^{-4}}{4}$       (B)  $-4e^{-4}$       (C)  $e^{-4} - 1$       (D)  $\frac{1}{4} - \frac{e^{-4}}{4}$       (E)  $4 - 4e^{-4}$

3. For  $x \geq 0$ , the horizontal line  $y = 2$  is an asymptote for the graph of the function  $f$ . Which of the following statements must be true?

- (A)  $f(0) = 2$   
 (B)  $f(x) \neq 2$  for all  $x \geq 0$   
 (C)  $f(2)$  is undefined.  
 (D)  $\lim_{x \rightarrow 2} f(x) = \infty$   
 (E)  $\lim_{x \rightarrow \infty} f(x) = 2$

6.  $\lim_{x \rightarrow \infty} \frac{x^3 - 2x^2 + 3x - 4}{4x^3 - 3x^2 + 2x - 1} =$

- (A) 4      (B) 1      (C)  $\frac{1}{4}$       (D) 0      (E) -1

4. If  $y = \frac{2x + 3}{3x + 2}$ , then  $\frac{dy}{dx} =$


- (A)  $\frac{12x + 13}{(3x + 2)^2}$       (B)  $\frac{12x - 13}{(3x + 2)^2}$       (C)  $\frac{5}{(3x + 2)^2}$       (D)  $\frac{-5}{(3x + 2)^2}$       (E)  $\frac{2}{3}$

5.  $\int_0^{\frac{\pi}{4}} \sin x dx =$

- (A)  $-\frac{\sqrt{2}}{2}$       (B)  $\frac{\sqrt{2}}{2}$       (C)  $-\frac{\sqrt{2}}{2} - 1$       (D)  $-\frac{\sqrt{2}}{2} + 1$       (E)  $\frac{\sqrt{2}}{2} - 1$

7. The graph of  $f'$ , the derivative of the function  $f$ , is shown above. Which of the following statements is true about  $f$ ?

- (A)  $f$  is decreasing for  $-1 \leq x \leq 1$ .  
 (B)  $f$  is increasing for  $-2 \leq x \leq 0$ .  
 (C)  $f$  is increasing for  $1 \leq x \leq 2$ .  
 (D)  $f$  has a local minimum at  $x = 0$ .  
 (E)  $f$  is not differentiable at  $x = -1$  and  $x = 1$ .



Graph of  $f'$

8.  $\int x^2 \cos(x^3) dx =$

(A)  $-\frac{1}{3} \sin(x^3) + C$

(B)  $\frac{1}{3} \sin(x^3) + C$

(C)  $-\frac{x^3}{3} \sin(x^3) + C$

(D)  $\frac{x^3}{3} \sin(x^3) + C$

(E)  $\frac{x^3}{3} \sin\left(\frac{x^4}{4}\right) + C$

9. If  $f(x) = \ln(x + 4 + e^{-3x})$ , then  $f'(0)$  is

(A)  $-\frac{2}{5}$

(B)  $\frac{1}{5}$


(C)  $\frac{1}{4}$

(D)  $\frac{2}{5}$


(E) nonexistent

10. The function  $f$  has the property that  $f(x)$ ,  $f'(x)$ , and  $f''(x)$  are negative for all real values  $x$ . Which of the following could be the graph of  $f$ ?


(A)




(B)




(C)



(D)



(E)



11. Using the substitution  $u = 2x + 1$ ,  $\int_0^2 \sqrt{2x + 1} dx$  is equivalent to


- (A)  $\frac{1}{2} \int_{-1/2}^{1/2} \sqrt{u} du$       (B)  $\frac{1}{2} \int_0^2 \sqrt{u} du$       (C)  $\frac{1}{2} \int_1^5 \sqrt{u} du$       (D)  $\int_0^2 \sqrt{u} du$       (E)  $\int_1^5 \sqrt{u} du$

12. The rate of change of the volume,  $V$ , of water in a tank with respect to time,  $t$ , is directly proportional to the square root of the volume. Which of the following is a differential equation that describes this relationship?

- (A)  $V(t) = k\sqrt{t}$   
 (B)  $V(t) = k\sqrt{V}$   
 (C)  $\frac{dV}{dt} = k\sqrt{t}$   
 (D)  $\frac{dV}{dt} = \frac{k}{\sqrt{V}}$   
 (E)  $\frac{dV}{dt} = k\sqrt{V}$

14. If  $y = x^2 \sin 2x$ , then  $\frac{dy}{dx} =$

- (A)  $2x \cos 2x$   
 (B)  $4x \cos 2x$   
 (C)  $2x(\sin 2x + \cos 2x)$   
 (D)  $2x(\sin 2x - x \cos 2x)$   
 (E)  $2x(\sin 2x + x \cos 2x)$



13. The graph of a function  $f$  is shown above. At which value of  $x$  is  $f$  continuous, but not differentiable?

- (A)  $a$       (B)  $b$       (C)  $c$       (D)  $d$       (E)  $e$

15. Let  $f$  be the function with derivative given by  $f'(x) = x^2 - \frac{2}{x}$ . On which of the following intervals is  $f$  decreasing?

- (A)  $(-\infty, -1]$  only  
 (B)  $(-\infty, 0)$   
 (C)  $[-1, 0)$  only  
 (D)  $(0, \sqrt[3]{2}]$   
 (E)  $[\sqrt[3]{2}, \infty)$

16. If the line tangent to the graph of the function  $f$  at the point  $(1, 7)$  passes through the point  $(-2, -2)$ , then  $f'(1)$  is

- (A)  $-5$       (B)  $1$       (C)  $3$       (D)  $7$       (E) undefined

17. Let  $f$  be the function given by  $f(x) = 2xe^x$ . The graph of  $f$  is concave down when

- (A)  $x < -2$       (B)  $x > -2$       (C)  $x < -1$       (D)  $x > -1$       (E)  $x < 0$

$x$	-4	-3	-2	-1	0	1	2	3	4
$g'(x)$	2	3	0	-3	-2	-1	0	3	2

18. The derivative  $g'$  of a function  $g$  is continuous and has exactly two zeros. Selected values of  $g'$  are given in the table above. If the domain of  $g$  is the set of all real numbers, then  $g$  is decreasing on which of the following intervals?

- (A)  $-2 \leq x \leq 2$  only  
 (B)  $-1 \leq x \leq 1$  only  
 (C)  $x \geq -2$   
 (D)  $x \geq 2$  only  
 (E)  $x \leq -2$  or  $x \geq 2$

19. A curve has slope  $2x + 3$  at each point  $(x, y)$  on the curve. Which of the following is an equation for this curve if it passes through the point  $(1, 2)$ ?


- (A)  $y = 5x - 3$   
 (B)  $y = x^2 + 1$   
 (C)  $y = x^2 + 3x$   
 (D)  $y = x^2 + 3x - 2$   
 (E)  $y = 2x^2 + 3x - 3$

20. Let  $f$  be the function given above. Which of the following statements are true about  $f$ ?

- I.  $\lim_{x \rightarrow 3} f(x)$  exists.  
 II.  $f$  is continuous at  $x = 3$ .  
 III.  $f$  is differentiable at  $x = 3$ .


$$f(x) = \begin{cases} x + 2 & \text{if } x \leq 3 \\ 4x - 7 & \text{if } x > 3 \end{cases}$$

- (A) None  
 (B) I only  
 (C) II only  
 (D) I and II only  
 (E) I, II, and III



21. The second derivative of the function  $f$  is given by  $f''(x) = x(x - a)(x - b)^2$ . The graph of  $f''$  is shown above. For what values of  $x$  does the graph of  $f$  have a point of inflection?


- (A) 0 and  $a$  only      (B) 0 and  $m$  only      (C)  $b$  and  $j$  only      (D) 0,  $a$ , and  $b$       (E)  $b$ ,  $j$ , and  $k$



22. The graph of  $f'$ , the derivative of  $f$ , is the line shown in the figure above. If  $f(0) = 5$ , then  $f(1) =$   
 (A) 0      (B) 3      (C) 6      (D) 8      (E) 11
23.  $\frac{d}{dx} \left( \int_0^{x^2} \sin(t^3) dt \right) =$   
 (A)  $-\cos(x^6)$       (B)  $\sin(x^3)$       (C)  $\sin(x^6)$       (D)  $2x \sin(x^3)$       (E)  $2x \sin(x^6)$
24. Let  $f$  be the function defined by  $f(x) = 4x^3 - 5x + 3$ . Which of the following is an equation of the line tangent to the graph of  $f$  at the point where  $x = -1$ ?  
 (A)  $y = 7x - 3$   
 (B)  $y = 7x + 7$   
 (C)  $y = 7x + 11$   
 (D)  $y = -5x - 1$   
 (E)  $y = -5x - 5$
25. A particle moves along the  $x$ -axis so that at time  $t \geq 0$  its position is given by  $x(t) = 2t^3 - 21t^2 + 72t - 53$ . At what time  $t$  is the particle at rest?  
 (A)  $t = 1$  only  
 (B)  $t = 3$  only  
 (C)  $t = \frac{7}{2}$  only  
 (D)  $t = 3$  and  $t = \frac{7}{2}$   
 (E)  $t = 3$  and  $t = 4$
26. What is the slope of the line tangent to the curve  $3y^2 - 2x^2 = 6 - 2xy$  at the point  $(3, 2)$ ?  
 (A) 0      (B)  $\frac{4}{9}$       (C)  $\frac{7}{9}$       (D)  $\frac{6}{7}$       (E)  $\frac{5}{3}$
27. Let  $f$  be the function defined by  $f(x) = x^3 + x$ . If  $g(x) = f^{-1}(x)$  and  $g(2) = 1$ , what is the value of  $g'(2)$ ?  
 (A)  $\frac{1}{13}$       (B)  $\frac{1}{4}$       (C)  $\frac{7}{4}$       (D) 4      (E) 13
28. Let  $g$  be a twice-differentiable function with  $g'(x) > 0$  and  $g''(x) > 0$  for all real numbers  $x$ , such that  $g(4) = 12$  and  $g(5) = 18$ . Of the following, which is a possible value for  $g(6)$ ?  
 (A) 15      (B) 18      (C) 21      (D) 24      (E) 27

76. A particle moves along the  $x$ -axis so that at any time  $t \geq 0$ , its velocity is given by  $v(t) = 3 + 4.1 \cos(0.9t)$ . What is the acceleration of the particle at time  $t = 4$ ?

- (A)  $-2.016$       (B)  $-0.677$       (C)  $1.633$       (D)  $1.814$       (E)  $2.978$






77. The regions  $A$ ,  $B$ , and  $C$  in the figure above are bounded by the graph of the function  $f$  and the  $x$ -axis. If the area of each region is 2, what is the value of  $\int_{-3}^3 (f(x) + 1) dx$ ?

- (A)  $-2$       (B)  $-1$       (C)  $4$       (D)  $7$       (E)  $12$

78. The radius of a circle is increasing at a constant rate of 0.2 meters per second. What is the rate of increase in the area of the circle at the instant when the circumference of the circle is  $20\pi$  meters?

- (A)  $0.04\pi \text{ m}^2/\text{sec}$   
 (B)  $0.4\pi \text{ m}^2/\text{sec}$   
 (C)  $4\pi \text{ m}^2/\text{sec}$   
 (D)  $20\pi \text{ m}^2/\text{sec}$   
 (E)  $100\pi \text{ m}^2/\text{sec}$

79. For which of the following does  $\lim_{x \rightarrow 4} f(x)$  exist?



- (A) I only  
 (B) II only  
 (C) III only  
 (D) I and II only  
 (E) I and III only

80. The function  $f$  is continuous for  $-2 \leq x \leq 1$  and differentiable for  $-2 < x < 1$ . If  $f(-2) = -5$  and  $f(1) = 4$ , which of the following statements could be false?

- (A) There exists  $c$ , where  $-2 < c < 1$ , such that  $f(c) = 0$ .  
 (B) There exists  $c$ , where  $-2 < c < 1$ , such that  $f'(c) = 0$ .  
 (C) There exists  $c$ , where  $-2 < c < 1$ , such that  $f(c) = 3$ .  
 (D) There exists  $c$ , where  $-2 < c < 1$ , such that  $f'(c) = 3$ .  
 (E) There exists  $c$ , where  $-2 \leq c \leq 1$ , such that  $f(c) \geq f(x)$  for all  $x$  on the closed interval  $-2 \leq x \leq 1$ .

81. Let  $f$  be the function with derivative given by  $f'(x) = \sin(x^2 + 1)$ . How many relative extrema does  $f$  have on the interval  $2 < x < 4$ ?

- (A) One      (B) Two      (C) Three      (D) Four      (E) Five

82. The rate of change of the altitude of a hot-air balloon is given by  $r(t) = t^3 - 4t^2 + 6$  for  $0 \leq t \leq 8$ . Which of the following expressions gives the change in altitude of the balloon during the time the altitude is decreasing?

(A)  $\int_{1.572}^{3.514} r(t) dt$

(B)  $\int_0^8 r(t) dt$

(C)  $\int_0^{2.667} r(t) dt$

(D)  $\int_{1.572}^{3.514} r'(t) dt$

(E)  $\int_0^{2.667} r'(t) dt$

83. The velocity, in ft/sec, of a particle moving along the  $x$ -axis is given by the function  $v(t) = e^t + te^t$ . What is the average velocity of the particle from time  $t = 0$  to time  $t = 3$ ?

(A) 20.086 ft/sec

(B) 26.447 ft/sec

(C) 32.809 ft/sec






(D) 40.671 ft/sec

(E) 79.342 ft/sec

84. A pizza, heated to a temperature of 350 degrees Fahrenheit ( $^{\circ}\text{F}$ ), is taken out of an oven and placed in a  $75^{\circ}\text{F}$  room at time  $t = 0$  minutes. The temperature of the pizza is changing at a rate of  $-110e^{-0.4t}$  degrees Fahrenheit per minute. To the nearest degree, what is the temperature of the pizza at time  $t = 5$  minutes?

- (A)  $112^{\circ}\text{F}$       (B)  $119^{\circ}\text{F}$       (C)  $147^{\circ}\text{F}$       (D)  $238^{\circ}\text{F}$       (E)  $335^{\circ}\text{F}$

85. If a trapezoidal sum overapproximates  $\int_0^4 f(x) dx$ , and a right Riemann sum underapproximates  $\int_0^4 f(x) dx$ , which of the following could be the graph of  $y = f(x)$ ?



86. The base of a solid is the region in the first quadrant bounded by the  $y$ -axis, the graph of  $y = \tan^{-1} x$ , the horizontal line  $y = 3$ , and the vertical line  $x = 1$ . For this solid, each cross section perpendicular to the  $x$ -axis is a square. What is the volume of the solid?






- (A) 2.561      (B) 6.612      (C) 8.046      (D) 8.755      (E) 20.773

87. The function  $f$  has first derivative given by  $f'(x) = \frac{\sqrt{x}}{1+x+x^3}$ . What is the  $x$ -coordinate of the inflection point of the graph of  $f$ ?

- (A) 1.008      (B) 0.473      (C) 0      (D) -0.278      (E) The graph of  $f$  has no inflection point.

88. On the closed interval  $[2, 4]$ , which of the following could be the graph of a function  $f$  with the property that

$$\frac{1}{4-2} \int_2^4 f(t) dt = 1?$$



89. Let  $f$  be a differentiable function with  $f(2) = 3$  and  $f'(2) = -5$ , and let  $g$  be the function defined by  $g(x) = xf(x)$ . Which of the following is an equation of the line tangent to the graph of  $g$  at the point where  $x = 2$ ?

- (A)  $y = 3x$   
 (B)  $y - 3 = -5(x - 2)$   
 (C)  $y - 6 = -5(x - 2)$   
 (D)  $y - 6 = -7(x - 2)$   
 (E)  $y - 6 = -10(x - 2)$

90. For all  $x$  in the closed interval  $[2, 5]$ , the function  $f$  has a positive first derivative and a negative second derivative. Which of the following could be a table of values for  $f$ ?

(A) 

$x$	$f(x)$
2	7
3	9
4	12
5	16

(B) 

$x$	$f(x)$
2	7
3	11
4	14
5	16

(C) 

$x$	$f(x)$
2	16
3	12
4	9
5	7

(D) 

$x$	$f(x)$
2	16
3	14
4	11
5	7

(E) 


$x$	$f(x)$
2	16
3	13
4	10
5	7



91. A particle moves along the  $x$ -axis so that at any time  $t > 0$ , its acceleration is given by  $a(t) = \ln(1 + 2^t)$ . If the velocity of the particle is 2 at time  $t = 1$ , then the velocity of the particle at time  $t = 2$  is
- (A) 0.462      (B) 1.609      (C) 2.555      (D) 2.886      (E) 3.346

92. Let  $g$  be the function given by  $g(x) = \int_0^x \sin(t^2) dt$  for  $-1 \leq x \leq 3$ . On which of the following intervals is  $g$  decreasing?

- (A)  $-1 \leq x \leq 0$   
 (B)  $0 \leq x \leq 1.772$   
 (C)  $1.253 \leq x \leq 2.171$   
 (D)  $1.772 \leq x \leq 2.507$   
 (E)  $2.802 \leq x \leq 3$




FR 1-3 w/ calc  
 4-6 non calc

- Let  $R$  be the shaded region bounded by the graphs of  $y = \sqrt{x}$  and  $y = e^{-3x}$  and the vertical line  $x = 1$ , as shown in the figure above.
  - Find the area of  $R$ .
  - Find the volume of the solid generated when  $R$  is revolved about the horizontal line  $y = 1$ .
  - The region  $R$  is the base of a solid. For this solid, each cross section perpendicular to the  $x$ -axis is a rectangle whose height is 5 times the length of its base in region  $R$ . Find the volume of this solid.
- A particle moves along the  $x$ -axis so that its velocity at time  $t$  is given by

$$v(t) = -(t + 1) \sin\left(\frac{t^2}{2}\right).$$


At time  $t = 0$ , the particle is at position  $x = 1$ .


- Find the acceleration of the particle at time  $t = 2$ . Is the speed of the particle increasing at  $t = 2$ ? Why or why not?
- Find all times  $t$  in the open interval  $0 < t < 3$  when the particle changes direction. Justify your answer.
- Find the total distance traveled by the particle from time  $t = 0$  until time  $t = 3$ .
- During the time interval  $0 \leq t \leq 3$ , what is the greatest distance between the particle and the origin? Show the work that leads to your answer.



$t$ (minutes)	$R(t)$ (gallons per minute)
0	20
30	30
40	40
50	55
70	65
90	70

3. The rate of fuel consumption, in gallons per minute, recorded during an airplane flight is given by a twice-differentiable and strictly increasing function  $R$  of time  $t$ . The graph of  $R$  and a table of selected values of  $R(t)$ , for the time interval  $0 \leq t \leq 90$  minutes, are shown above.
- Use data from the table to find an approximation for  $R'(45)$ . Show the computations that lead to your answer. Indicate units of measure.
  - The rate of fuel consumption is increasing fastest at time  $t = 45$  minutes. What is the value of  $R''(45)$ ? Explain your reasoning.
  - Approximate the value of  $\int_0^{90} R(t) dt$  using a left Riemann sum with the five subintervals indicated by the data in the table. Is this numerical approximation less than the value of  $\int_0^{90} R(t) dt$ ? Explain your reasoning.
  - For  $0 < b \leq 90$  minutes, explain the meaning of  $\int_0^b R(t) dt$  in terms of fuel consumption for the plane. Explain the meaning of  $\frac{1}{b} \int_0^b R(t) dt$  in terms of fuel consumption for the plane. Indicate units of measure in both answers.
4. Let  $f$  be a function defined on the closed interval  $-3 \leq x \leq 4$  with  $f(0) = 3$ . The graph of  $f'$ , the derivative of  $f$ , consists of one line segment and a semicircle, as shown above.
- On what intervals, if any, is  $f$  increasing? Justify your answer.
  - Find the  $x$ -coordinate of each point of inflection of the graph of  $f$  on the open interval  $-3 < x < 4$ . Justify your answer.
  - Find an equation for the line tangent to the graph of  $f$  at the point  $(0, 3)$ .
  - Find  $f(-3)$  and  $f(4)$ . Show the work that leads to your answers.





5. A coffee pot has the shape of a cylinder with radius 5 inches, as shown in the figure above. Let  $h$  be the depth of the coffee in the pot, measured in inches, where  $h$  is a function of time  $t$ , measured in seconds. The volume  $V$  of coffee in the pot is changing at the rate of  $-5\pi\sqrt{h}$  cubic inches per second. (The volume  $V$  of a cylinder with radius  $r$  and height  $h$  is  $V = \pi r^2 h$ .)

- (a) Show that  $\frac{dh}{dt} = -\frac{\sqrt{h}}{5}$ .
- (b) Given that  $h = 17$  at time  $t = 0$ , solve the differential equation  $\frac{dh}{dt} = -\frac{\sqrt{h}}{5}$  for  $h$  as a function of  $t$ .
- (c) At what time  $t$  is the coffee pot empty?

6. Let  $f$  be the function defined by

$$f(x) = \begin{cases} \sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ 5-x & \text{for } 3 < x \leq 5. \end{cases}$$

- (a) Is  $f$  continuous at  $x = 3$ ? Explain why or why not.
- (b) Find the average value of  $f(x)$  on the closed interval  $0 \leq x \leq 5$ .
- (c) Suppose the function  $g$  is defined by

$$g(x) = \begin{cases} k\sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ mx+2 & \text{for } 3 < x \leq 5, \end{cases}$$

where  $k$  and  $m$  are constants. If  $g$  is differentiable at  $x = 3$ , what are the values of  $k$  and  $m$ ?